



Neck Circumference and Insulin Resistance in Chinese Adults: The Cardiometabolic Risk in Chinese (CRC) Study

Citation

Liang, Jun, Fei Teng, Yanping Li, Xuekui Liu, Caiyan Zou, Yu Wang, Hongyan Li, and Lu Qi. 2013. "Neck Circumference and Insulin Resistance in Chinese Adults: The Cardiometabolic Risk in Chinese (CRC) Study." *Diabetes Care* 36 (9): e145-e146. doi:10.2337/dc13-1114. <http://dx.doi.org/10.2337/dc13-1114>.

Published Version

doi:10.2337/dc13-1114

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:12987237>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

OBSERVATIONS

Neck Circumference and Insulin Resistance in Chinese Adults: The Cardiometabolic Risk in Chinese (CRC) Study

Neck circumference (NC) is a proxy for upper-body fat and a reliable screening measure for the identification of individuals with abnormal fat distribution. Recently, compelling evidence indicates that NC is positively associated with insulin resistance (IR) and a variety of cardiometabolic abnormalities, independent of overall adiposity and central obesity in Caucasians (1) or Chinese diabetic patients (2). However, it remains unknown whether NC affects IR in the general Chinese population.


In the current study, we analyzed the associations of NC with IR and cardiometabolic risk factors in 2,588 apparently healthy Chinese adults (20–87 years of age) from the Cardiometabolic Risk in Chinese (CRC) study (3). We found a significant association between NC and an increasing trend of homeostasis model assessment of insulin resistance (HOMA-IR), adjusting for age and other cardiometabolic risk factors, in both men ($P = 0.02$) and women ($P = 0.004$). The differences in HOMA-IR between the extreme quintiles of NC appeared more evident in men than in women; however, the test for the heterogeneity between men and women was not significant. Further adjustment for BMI and waist circumference (WC) did not appreciably change the associations between NC and IR. In addition, high NC was significantly related to a decreasing trend of HDL levels in both sexes and was significantly related to an increasing trend of plasma triglyceride and uric acid levels only in men (P for sex difference <0.03), after adjustment for covariates including BMI or WC and HOMA-IR.

We also found significant interactions of NC with BMI and WC (P for interaction <0.0001) in relation to HOMA-IR (Fig. 1). It appears that the associations between NC and HOMA-IR were more evident in those with a higher BMI or WC.

To our knowledge, the current study is among the first to assess the associations of NC with IR and metabolic risk in a general population of Chinese adults. Our findings are in line with the results from several previous studies in Caucasians and Chinese diabetic patients (1,2). In the Framingham Study, it was found that NC was associated with IR and cardiovascular risk factors in a sex-specific manner (1). It has been documented that high NC was a significant predictor of obstructive sleep apnea syndrome (4), which has been associated with risk of IR and diabetes. Obstructive sleep apnea syndrome aggravates glycemic control, even at the earliest stages of glucose intolerance. In addition, intermittent hypoxemia

and sleep fragmentation increase the risk of IR (5). Our data, together with evidence from other studies, suggest that body fat accumulated in the upper body segment may contribute to the adverse metabolic consequence.


In conclusion, we found significant associations of high NC with increased risk of IR and a variety of cardiometabolic risk factors in apparently healthy Chinese adults, and found sexual dimorphism for associations with certain risk factors. Our findings support a recommendation for the inclusion of NC in the routine clinical assessment of adults to identify those at high risk of cardiometabolic diseases such as type 2 diabetes and heart disease.



Low: WC < 74 cm for women, WC < 86 cm for men; P for trend = 0.0003

Medium: $74\text{cm} \leq \text{WC} < 81\text{cm}$ for women, $86\text{cm} \leq \text{WC} < 93\text{cm}$ for men; P for trend = 0.0023

High: WC $\geq 81\text{cm}$ for women, WC $\geq 93\text{cm}$ for men; P for trend < 0.0001



Low: $22.99 < \text{BMI}$; P for trend = 0.5948

Medium: $22.99 \leq \text{BMI} < 25.65$; P for trend = 0.0004

High: BMI ≥ 25.65 ; P for trend < 0.0001

Figure 1—The joint effect of NC (in quintiles) with WC and BMI (low, medium, and high levels) on HOMA-IR.

JUN LIANG, MD, PHD¹
 FEI TENG, MS¹
 YANPING LI, MD, PHD²
 XUEKUI LIU, BS¹
 CAIYAN ZOU, MS¹
 YU WANG, MS¹
 HONGYAN LI, MS¹
 LU QI, MD, PHD^{2,3}

From the ¹Department of Endocrinology, Xuzhou Central Hospital, Xuzhou Clinical School of Xuzhou Medical College, Affiliated Hospital of Southeast University, Xuzhou, Jiangsu, China; the ²Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts; and the ³Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts.

Corresponding author: Lu Qi, nhlqi@channing.harvard.edu, or Jun Liang, mwlj521@163.com.

J.L. and F.T. contributed equally to this study.

DOI: 10.2337/dc13-1114

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—L.Q.'s research is supported by National Institutes of Health grants DK-091718 and HL-071981, the American Heart Association Scientist Development Award, and the Boston Obesity Nutrition Research Center (DK-46200). J.L.'s research was sponsored by Jiangsu Health International Exchange Program and supported by a Xuzhou medical leading talent grant and Xuzhou Science and Technology Board Foundation grant (XF11C067).

No potential conflicts of interest relevant to this article were reported.

J.L. wrote and reviewed the manuscript. F.T. wrote the manuscript. Y.L. and X.L. performed statistical analyses. C.Z. and Y.W. contributed to discussion. H.L. edited the manuscript. L.Q. researched data and wrote the manuscript. L.Q. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The authors thank all subjects for participating in this study.

References

1. Preis SR, Massaro JM, Hoffmann U, et al. Neck circumference as a novel measure of cardio-metabolic risk: the Framingham Heart study. *J Clin Endocrinol Metab* 2010;95:3701–3710
2. Yang GR, Yuan SY, Fu HJ, et al.; Beijing Community Diabetes Study Group. Neck circumference positively related with central obesity, overweight, and metabolic syndrome in Chinese subjects with type 2 diabetes: Beijing Community Diabetes Study 4. *Diabetes Care* 2010;33:2465–2467
3. Liang J, Zhou N, Teng F, et al. Hemoglobin A1c levels and aortic arterial stiffness: the Cardiometabolic Risk in Chinese (CRC) study. *PLoS ONE* 2012;7:e38485
4. Onat A, Hergenç G, Yüksel H, et al. Neck circumference as a measure of central obesity: associations with metabolic syndrome and obstructive sleep apnea syndrome beyond waist circumference. *Clin Nutr* 2009;28:46–51
5. Steiropoulos P, Papanas N, Bouros D, Maltezos E. Obstructive sleep apnea aggravates glycemic control across the continuum of glucose homeostasis. *Am J Respir Crit Care Med* 2010;182:286